

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
19 February 2004 (19.02.2004)

PCT

(10) International Publication Number  
**WO 2004/014373 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/436**,  
A61P 27/02

(21) International Application Number:  
PCT/JP2003/010105

(22) International Filing Date: 8 August 2003 (08.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/402,061 9 August 2002 (09.08.2002) US

(71) Applicant (*for all designated States except US*): **SU-  
CAMPO PHARMACEUTICALS, INC.** [US/US]; 4733  
Bethesda Avenue, Suite 450, Bethesda, MD 20814 (US).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **UENO, Ryuji** [JP/US];  
11025 Stanmore Drive, Potomac, Montgomery, MD 20854  
(US).

(74) Agent: **TAKASHIMA, Hajime**; Fujimura Yamato Seimei  
Bldg., 2-14, Fushimimachi 4-chome, Chuo-ku, Osaka-shi,  
Osaka 541-0044 (JP).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD,  
SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,  
US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PHARMACEUTICAL COMPOSITIONS COMPRISING FK506 DERIVATIVES AND THE IR USE FOR THE TREATMENT OF ALLERGIC DISEASES

(57) Abstract: The invention provides ophthalmic compositions and methods treating the symptoms of ocular allergies. The principle active ingredient in these compositions and methods is macrolide compound, such as tacrolimus, ascomycin and rapamycin and their derivatives. Optimal concentrations and dosing regimens are provided.

WO 2004/014373 A1

## Description

PHARMACEUTICAL COMPOSITIONS COMPRISING FK506 DERIVATIVES AND THEIR USE FOR THE TREATMENT OF ALLERGIC DISEASES

### Technical Field of the Invention

5           The present invention relates to a method for treating ocular allergies.

### Background Art

          The incidence and prevalence of allergic conjunctivitis has increased dramatically over the past 40 years, today affecting up to 20% of the US population. The condition can be seasonal if due to pollens from trees, grasses or weeds, or  
10   perennial, if the antigen is abundant throughout the year such as animal dander, dust or mold; of the two, seasonal allergic conjunctivitis is more common. Ophthalmologists believe that 80%-90% of all allergic conjunctivitis cases are seasonal, while the remaining 10%-20% are perennial in nature.

          Ocular allergies, like allergic conjunctivitis are currently being served with  
15   products within the following categories: antihistamines, mast cell stabilizers, nonsteroidal anti-inflammatory drugs ("NSAIDS") and corticosteroids. Despite the availability of so many products, none of them are entirely satisfactory and there still exists a need for products that are effective and work using different mechanisms of action.

20           Historically, there has been some interest in the development of macrolide immunosuppressive compounds in to treat allergic conjunctivitis, but as yet there is no product on the market. Of these macrolide compounds, notable is tacrolimus, aka FK506, originated by Fujisawa Pharmaceutical Co., Ltd. in Japan. See U.S. Patent No. 5,514,686. Heretofore, however, there has never been defined an optimal dosing  
25   regimen for treating human patients.

### Disclosure of the Invention

          It is, therefore, an object of the invention to provide useful, improved compositions and methods for treating ocular allergy symptoms with macrolide immunosuppressive compounds.


30           An another object of the invention to provide a commercial package comprising the composition of the present invention and a written matter associated

therewith, the written matter stating the composition can or should be used for ocular allergies, especially allergic conjunctivitis.

According to this and other objects of the invention, a method of treating a human patient suffering from ocular allergy symptoms is provided. According to one embodiment, this method entails administering to the patient an ophthalmic composition containing from about 0.01% to about 0.1% of a macrolide compound. In other embodiments, the method involves administering to the patient an ophthalmic composition containing from about 0.03% to about 0.06% of a macrolide compound, but preferably about 0.03%.

Preferred compositions are formulated as eye drops, which optionally contain polyvinyl alcohol, or ointments. In general, these compositions will be administered to the eye from about one to about four times per day.

Preferred macrolide compound is a tricyclo compound having the following formula (I) or a pharmaceutically acceptable salt thereof:



wherein adjacent pairs of  $R^1$  and  $R^2$ ,  $R^3$  and  $R^4$ , and  $R^5$  and  $R^6$  each independently: (a) consist of two adjacent hydrogen atoms, wherein  $R^2$  is optionally alkyl, or (b) form another bond optionally between carbon atoms binding with the members of said pairs;  $R^7$  is hydrogen atom, hydroxy, alkyloxy or protected hydroxy, or may form oxo with  $R^1$ ;  $R^8$  and  $R^9$  each independently show hydrogen atom or hydroxy;  $R^{10}$  is hydrogen atom, alkyl, alkyl substituted by one or more hydroxy, alkenyl, alkenyl substituted by one or more hydroxy or alkyl substituted by oxo; X is oxo, (hydrogen

atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $-\text{CH}_2\text{O}-$ ; Y is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $\text{N}-\text{NR}^{11}\text{R}^{12}$  or  $\text{N}-\text{OR}^{13}$ ;  $\text{R}^{11}$  and  $\text{R}^{12}$  each independently show hydrogen atom, alkyl, aryl or tosyl;  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{18}$ ,  $\text{R}^{19}$ ,  $\text{R}^{22}$  and  $\text{R}^{23}$  each

- 5 independently show hydrogen atom or alkyl;  $\text{R}^{24}$  is an optionally substituted ring that may contain one or more hetero atom(s); and n is 1 or 2. Tacrolimus is most preferred.

### Brief Description of the Drawing

Figure 1 shows the ability of macrolide-containing eye drops to suppress ocular itching in humans in response to challenge with common allergens.


10

### Detailed Description of the Invention

The present inventor has surprisingly discovered that certain macrolide compounds can be used in specific concentration ranges to treat the ocular symptoms of allergy. In particular, macrolide compounds like FK506 (tacrolimus), ascomycin, rapamycin and their derivatives, can be used in concentrations ranging from about 15 0.01% to about 0.1% in ophthalmic compositions to treat ocular allergy symptoms and, in particular, allergic conjunctivitis.

### Macrolide Compounds of the Invention

A specific example of a macrolide compound usable in the invention is a tricyclo compound as shown by the following general formula (I) or a 20 pharmaceutically acceptable salt thereof.



wherein adjacent pairs of  $\text{R}^1$  and  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$ , and  $\text{R}^5$  and  $\text{R}^6$  each independently

a) consist of two adjacent hydrogen atoms, wherein  $R^2$  is optionally alkyl, or

b) form another bond optionally between carbon atoms binding with the members of said pairs;

$R^7$  is hydrogen atom, hydroxy, alkyloxy or protected hydroxy, or may form  
5 oxo with  $R^1$ ;

$R^8$  and  $R^9$  each independently show hydrogen atom or hydroxy;

$R^{10}$  is hydrogen atom, alkyl, alkyl substituted by one or more hydroxy, alkenyl, alkenyl substituted by one or more hydroxy or alkyl substituted by oxo;

X is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a  
10 group of the formula  $-CH_2O-$ ;

Y is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $N-NR^{11}R^{12}$  or  $N-OR^{13}$ ;

$R^{11}$  and  $R^{12}$  each independently show hydrogen atom, alkyl, aryl or tosyl;

$R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$ ,  $R^{19}$ ,  $R^{22}$  and  $R^{23}$  each independently show  
15 hydrogen atom or alkyl;

$R^{24}$  is an optionally substituted ring that may contain one or more hetero atom(s) and;

n is 1 or 2.

In addition to the meaning noted above, Y,  $R^{10}$  and  $R^{23}$  may show, together  
20 with the carbon atom they bind with, a saturated or unsaturated 5 or 6-membered heterocyclic group containing nitrogen atom, sulfur atom and/or oxygen atom, the heterocyclic group being optionally substituted by one or more group(s) selected from alkyl, hydroxy, alkyloxy, benzyl, a group of the formula  $-CH_2Se(C_6H_5)$ , and alkyl substituted by one or more hydroxy, or its pharmaceutically acceptable salt.

25 In the general formula (I), preferably  $R^{24}$  is, for example, cyclo( $C_5-C_7$ )alkyl optionally having suitable substituent, such as the following.

(a) 3,4-dioxocyclohexyl

(b) 3- $R^{20}$ -4- $R^{21}$ -cyclohexyl,

wherein R<sup>20</sup> is hydroxy, alkyloxy or -OCH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, and R<sup>21</sup> is hydroxy, -OCN, alkyloxy, heteroaryloxy optionally having suitable substituent, -OCH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, protected hydroxy, chloro, bromo, iodo, aminooxalyloxy, azide, p-tolyloxythiocarbonyloxy, or R<sup>25</sup>R<sup>26</sup>CHCOO-(wherein R<sup>25</sup> is hydroxy optionally protected where desired or protected amino, and R<sup>26</sup> is hydrogen atom or methyl, or R<sup>20</sup> and R<sup>21</sup> in combination form an oxygen atom of epoxide ring or

(c) cyclopentyl wherein cyclopentyl is substituted by methoxymethyl, optionally protected hydroxymethyl where desired, acyloxymethyl (wherein acyl moiety is optionally quaternized dimethylamino or optionally esterified carboxy), one or more optionally protected amino and/or hydroxy, or aminooxalyloxymethyl. Preferable examples include 2-formyl-cyclopentyl.

The definition of each symbol used in the formula (I), specific examples thereof and preferable embodiments thereof will be explained in detail in the following.

“Lower” generally means a group having from about 1 to about 6 carbon atoms unless otherwise indicated.

Preferable examples of the alkyl moiety of “alkyl” and “alkyloxy” include linear or branched aliphatic hydrocarbon residue, such as lower alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, neopentyl, hexyl and the like).

Preferable examples of “alkenyl” include linear or branched aliphatic hydrocarbon residue having one double bond, such as lower alkenyl (e.g., vinyl, propenyl (e.g., allyl and the like), butenyl, methylpropenyl, pentenyl, hexenyl and the like).

Preferable examples of “aryl” include phenyl, tolyl, xylyl, cumenyl, mesityl, naphthyl and the like.

Preferable examples of the protective group for “protected hydroxy” and “protected amino” include 1-(lower alkylthio)(lower)alkyl such as lower alkylthiomethyl (e.g., methylthiomethyl, ethylthiomethyl, propylthiomethyl, isopropylthiomethyl, butylthiomethyl, isobutylthiomethyl, hexylthiomethyl and the like), with more preference given to C<sub>1</sub> - C<sub>4</sub> alkylthiomethyl and most preference given to methylthiomethyl; tri-substituted silyl such as tri(lower)alkylsilyl (e.g.,

trimethylsilyl, triethylsilyl, tributylsilyl, tert-butyl dimethylsilyl, tri-tert-butylsilyl and the like), and lower alkyldiarylsilyl (e.g., methyldiphenylsilyl, ethyldiphenylsilyl, propyldiphenylsilyl, tert-butyldiphenylsilyl and the like), with more preference given to tri(C<sub>1</sub> - C<sub>4</sub>)alkylsilyl and C<sub>1</sub> - C<sub>4</sub> alkyldiphenylsilyl, and most preference given to  
5 tert-butyl-dimethylsilyl and tert-butyldiphenylsilyl; acyl such as aliphatic acyl, aromatic acyl and aliphatic acyl substituted by aromatic group, which are derived from carboxylic acid, sulfonic acid and carbamic acid; and the like.

The aliphatic acyl is exemplified by lower alkanoyl optionally having one or more suitable substituent(s) (e.g., carboxy) such as formyl, acetyl, propionyl, butyryl,  
10 isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, carboxyacetyl, carboxypropionyl, carboxybutyryl, carboxyhexanoyl and the like; cyclo(lower)alkyloxy(lower)alkanoyl optionally having one or more suitable substituent(s) (e.g., lower alkyl) such as cyclopropyloxyacetyl, cyclobutyloxypropionyl, cycloheptyloxybutyryl, mentyloxyacetyl, mentyloxypropionyl, mentyloxybutyryl, mentyloxypentanoyl,  
15 mentyloxyhexanoyl and the like; camphorsulfonyl; lower alkylcarbamoyl having one or more suitable substituent(s) such as carboxy or protected carboxy and the like, such as carboxy(lower)alkylcarbamoyl (e.g., carboxymethylcarbamoyl, carboxyethylcarbamoyl, carboxypropylcarbamoyl, carboxybutylcarbamoyl, carboxypentylcarbamoyl, carboxyhexylcarbamoyl) and  
20 tri(lower)alkylsilyl(lower)alkyloxycarbonyl(lower)alkylcarbamoyl (e.g., trimethylsilylmethoxycarbonylethylcarbamoyl, trimethylsilylethoxycarbonylpropylcarbamoyl, triethylsilylethoxycarbonylpropylcarbamoyl, tert-butyl dimethylsilylethoxycarbonylpropylcarbamoyl,  
25 trimethylsilylpropoxycarbonylbutylcarbamoyl.

Aromatic acyl is exemplified by aroyl optionally having one or more suitable substituent(s) (e.g., nitro), such as benzoyl, toluoyl, xyloyl, naphthoyl, nitrobenzoyl, dinitrobenzoyl, nitronaphthoyl and the like and arenesulfonyl optionally having one or more suitable substituent(s) (e.g., halogen), such as benzenesulfonyl, toluenesulfonyl,  
30 xylenesulfonyl, naphthalenesulfonyl, fluorobenzenesulfonyl, chlorobenzenesulfonyl, bromobenzenesulfonyl, iodobenzenesulfonyl and the like.

The aliphatic acyl substituted by aromatic group may be, for example, ar(lower)alkanoyl optionally having one or more suitable substituent(s) (e.g., lower alkyloxy or trihalo(lower)alkyl and the like), wherein specific examples are phenylacetyl, phenylpropionyl, phenylbutyryl, 2-trifluoromethyl-2-methoxy-2-phenylacetyl, 2-ethyl-2-trifluoromethyl-2-phenylacetyl, 2-trifluoromethyl-2-propoxy-2-phenylacetyl and the like.

Of the above-mentioned acyl, more preferable acyl includes C<sub>1</sub> - C<sub>4</sub> alkanoyl optionally having carboxy, cyclo(C<sub>5</sub> - C<sub>6</sub>)alkyloxy(C<sub>1</sub> - C<sub>4</sub>)alkanoyl having two (C<sub>1</sub> - C<sub>4</sub>)alkyl in the cycloalkyl moiety, camphorsulfonyl, carboxy (C<sub>1</sub> - C<sub>4</sub>)alkylcarbamoyl, tri(C<sub>1</sub> - C<sub>4</sub>)alkylsilyl(C<sub>1</sub> - C<sub>4</sub>)alkyloxycarbonyl(C<sub>1</sub> - C<sub>4</sub>)alkylcarbamoyl, benzoyl optionally having one or two nitro groups, and benzenesulfonyl having halogen, phenyl(C<sub>1</sub> - C<sub>4</sub>)alkanoyl having C<sub>1</sub> - C<sub>4</sub> alkyloxy and trihalo(C<sub>1</sub> - C<sub>4</sub>)alkyl. Of these, most preferred are acetyl, carboxypropionyl, mentyloxyacetyl, camphorsulfonyl, benzoyl, nitrobenzoyl, dinitrobenzoyl, iodobenzenesulfonyl, 2-trifluoromethyl-2-methoxy-2-phenylacetyl and the like.

Preferable examples of the "heterocyclic group consisting of saturated or unsaturated 5 or 6-membered ring having nitrogen atom, sulfur atom and/or oxygen atom" are pyrrolyl, tetrahydrofuryl and the like.

The "heteroaryl optionally having suitable substituent moiety" of the "heteroaryloxy optionally having suitable substituent" is that exemplified for R<sup>1</sup> of the compound of the formula I of EP-A-532,088, with preference given to 1-hydroxyethylindol-5-yl. The disclosure is incorporated herein by reference.


The tricyclo compound (I) used in the present invention is described in the publications EP-A-184162, EP-A-323042, EP-A-423714, EP-A-427680, EP-A-465426, EP-A-480623, EP-A-532088, EP-A-532089, EP-A-569337, EP-A-626385, WO89/05303, WO93/05058, WO96/31514, WO91/13889, WO91/19495, WO93/5059 and the like. The disclosures of these publications are incorporated herein by reference.

In particular, the compounds called FR900506 (FK506), FR900520 (Ascomycin), FR900523 and FR900525 are produced by the genus *Streptomyces*, such as *Streptomyces tsukubaensis*, No. 9993 (depository: National Institute of Advanced Industrial Science and Technology, International Patent Organism



Depository, Central 6, 1-1, Higashi 1-chome, Tsukuba-shi, Ibaraki-ken, Japan (formerly Fermentation Research Institute, Agency of Industrial Science and Technology, the Ministry of International Trade and Industry), date of deposit: October 5, 1984, deposit number FERM BP-927) or *Streptomyces hygroscopicus*

- 5 *subsp. Yakushimaensis*, No. 7238 (depository National Institute of Advanced Industrial Science and Technology, International Patent Organism Depository, Central 6, 1-1, Higashi 1-chome, Tsukuba-shi, Ibaraki-ken, Japan (formerly Fermentation Research Institute, Agency of Industrial Science and Technology, the Ministry of International Trade and Industry), date of deposit January 12, 1985, deposit number:
- 10 FERM BP-928 (EP-A-0184162)), and the compound of the following formula, FK506 (generic name: Tacrolimus) is a representative compound.



Chemical name: 17-allyl-1,14-dihydroxy-12-[2-(4-hydroxy-3-methoxycyclohexyl)-1-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.0<sup>4,9</sup>]octacos-18-ene-2,3,10,16-tetraone

- 15 Of the tricyclo compounds (I), more preferred is a compound wherein adjacent pairs of R<sup>3</sup> and R<sup>4</sup>, and R<sup>5</sup> and R<sup>6</sup> each independently form another bond optionally between carbon atoms binding with the members of said pairs;

R<sup>8</sup> and R<sup>23</sup> each independently show hydrogen atom;

R<sup>9</sup> is hydroxy;

- 20 R<sup>10</sup> is methyl, ethyl, propyl or allyl;

X is (hydrogen atom, hydrogen atom) or oxo;

Y is oxo;

R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup> and R<sup>22</sup> each independently show methyl;

R<sup>24</sup> is 3-R<sup>20</sup>-4-R<sup>21</sup>-cyclohexyl,

wherein  $R^{20}$  is hydroxy, alkyloxy or  $-OCH_2OCH_2CH_2OCH_3$ , and  $R^{21}$  is hydroxy,  $-OCN$ , alkyloxy, heteroaryloxy optionally having suitable substituent,  $-OCH_2OCH_2CH_2OCH_3$ , protected hydroxy, chloro, bromo, iodo, aminooxyloxy, azide, p-tolyloxythiocarbonyloxy or  $R^{25}R^{26}CHCOO-$  (wherein  $R^{25}$  is optionally protected hydroxy as desired, or protected amino, and  $R^{26}$  is hydrogen atom or methyl), or  $R^{20}$  and  $R^{21}$  in combination form an oxygen atom of epoxide ring; and  
n is 1 or 2.

Particularly preferable tricyclo macrolide compounds (I) include, besides FK506, Ascomycin derivatives such as halogenated derivative of 33-epi-chloro-33-desoxy Ascomycin described in Example 66a of EP-A-427,680 and the like.

Other preferable macrolide compounds include Rapamycin described in MERCK INDEX, 12 edition, No. 8288 and derivatives thereof. Preferable examples thereof include O-substituted derivative described at page 1 of WO95/16691, formula A, wherein the 40<sup>th</sup> hydroxy is  $-OR_1$  (wherein  $R_1$  is hydroxyalkyl, hydroalkyloxyalkyl, acylaminoalkyl and aminoalkyl), such as 40-O-(2-hydroxy)ethyl Rapamycin, 40-O-(3-hydroxy)propyl Rapamycin, 40-O-[2-(2-hydroxy)ethoxy]ethyl Rapamycin and 40-O-(2-acetaminoethyl)-Rapamycin. These O-substituted derivatives can be produced by reacting, under appropriate conditions, Rapamycin (or dihydro or deoxo Rapamycin) and an organic radical bound with leaving group (e.g.,  $RX$  wherein  $R$  is an organic radical desirable as O-substituent, such as alkyl, allyl and benzyl moiety, and  $X$  is a leaving group such as  $CCl_3C(NH)O$  and  $CF_3SO_3$ ). The conditions are: when  $X$  is  $CCl_3C(NH)O$ , acidic or neutral conditions, such as in the presence of trifluoromethanesulfonic acid, camphorsulfonic acid, p-toluenesulfonic acid or their corresponding pyridinium or substituted pyridinium salt, and when  $X$  is  $CF_3SO_3$ , in the presence of a base such as pyridine, substituted pyridine, diisopropylethylamine and pentamethylpiperidine. The most preferable Rapamycin derivative is 40-O-(2-hydroxy)ethyl Rapamycin as disclosed in WO94/09010, which is hereby incorporated into the specification by reference.

The pharmaceutically acceptable salt of tricyclo compound (I), Rapamycin and derivatives thereof are nontoxic and pharmaceutically acceptable conventional salts, which are exemplified by salts with inorganic or organic base such as alkali metal salt (e.g., sodium salt, potassium salt and the like), alkaline earth metal salt (e.g.,

calcium salt, magnesium salt and the like), ammonium salt, and amine salt (e.g., triethylamine salt, N-benzyl-N-methylamine salt and the like).

5 The macrolide compound of the invention comprises one or more pairs of stereoisomers, such as optical isomers and geometric isomers, which may be included due to conformers or asymmetric carbon atoms and double bonds. Such conformers and isomers are also encompassed in the present invention. In addition, macrolide compounds can form solvates, which also are encompassed by the present invention. Preferable solvates include hydrates and ethanolates.

10 The instant macrolide compounds and their pharmaceutically acceptable salts are nontoxic. Pharmaceutically acceptable conventional salts may have an inorganic or organic base, such as alkali metal salt (e.g., sodium salt, potassium salt and the like), alkaline earth metal salt (e.g., calcium salt, magnesium salt and the like), ammonium salt, and amine salt (e.g., triethylamine salt, N-benzyl-N-methylamine salt and the like).

15 As used herein, unless otherwise specifically noted, the term "macrolide" or reference to a particular macrolide is meant to include all pharmaceutically acceptable salts thereof.

### ***Ophthalmic Compositions***

20 While the present macrolide compounds may be administered any number of ways, the most convenient forms are contemplated to be eye drops and ointments, which may be prepared according to conventional methods. The optimal concentration of the macrolide compounds is in the range of about 0.01% to about 0.1% (more strictly, 0.01% to 0.1%), but more preferably is about 0.03% to about 0.06% (more strictly, 0.03% to 0.06%), with 0.03% being most preferred.

25 Eye drops, for instance, may be prepared by dissolving the active ingredient in a sterile aqueous solution such as physiological saline, buffering solution, etc., or by providing a powdered composition that is dissolved before use. Eye drops such as the ones as described in EP-A-0406791 (which is incorporated by reference in its entirety) are preferred. Conventional eye drop additives can be used. Such additives  
30 include isotonizing agents (e.g., sodium chloride, etc.), buffer agents (e.g., boric acid, sodium monohydrogen phosphate, sodium dihydrogen phosphate, etc.), preservatives (e.g., benzalkonium chloride, benzethonium chloride, chlorobutanol, etc.), thickeners

(e.g., saccharide such as lactose, mannitol, maltose, etc.; e.g., hyaluronic acid or its salt such as sodium hyaluronate, potassium hyaluronate, etc.; e.g., mucopolysaccharide such as chondroitin sulfate, etc.; e.g., sodium polyacrylate, carboxyvinyl polymer, crosslinked polyacrylate, etc.; e.g., polyvinyl alcohol, methylcellulose, glycerine, etc.).

Especially, polyvinyl alcohol as additive is preferably used in the eye drop of the present invention.

Ophthalmic ointments may be prepared by mixing the active ingredient with a base according to conventional methods. Examples ointment bases include, but are not limited to, petrolatum, selen 50, Plastibase and macrogol. In order to increase the hydrophilicity, a surface-active agent, like a detergent or other emulsifier, can be added. The same additives used in the eye drops, such as the preservatives, etc. can also be used in an ointment.

The present formulation can further include other pharmacological active ingredients as far as they do not contradict the purpose of the present invention. For instance, the formulation can include a single or multiple macrolide compounds, and may also include one or more antimicrobial agents as active ingredients for the purpose of treating or preventing bacterial infections. In a combination of plural active ingredients, their respective contents may be suitably increased or decreased in consideration of their effects and safety.

The present agent can be formulated as a sterile unit dose type containing no preservatives.

### ***Methods of Treatment***

The term "treatment" used herein includes any means of control such as prevention, care, relief of the condition, attenuation of the condition and arrest of progression.

The patient being treated will generally have a history of ocular allergy symptoms. Most pronounced among those symptoms are redness and itching. The patient may be suffering from allergic conjunctivitis.

The present macrolide-containing compositions, described above, generally are topically administered to the eyes and/or the surrounding skin, such as the eyelids.

The amount and frequency of administration can vary according to sex, age and weight of a human, symptoms to be treated, desirable therapeutic effects, administration routes and period for treatment. However, the inventor has found the optimal concentration of macrolide compound in the ophthalmic composition (eye drop, eye ointment) for treating ocular allergies to be in the range of about 0.01% to about 0.06%. Concentrations of up to about 0.1% may be used, but generally those are best formulated as an ointment. When considering all factors, concentrations of 0.03% appear to be best suited for treatment. Preferably, the macrolide compounds is formulated as an eye drop and may be administered several times a day per eye, preferably one to six times, more preferably one to four times, several drops per time, preferably one to four drops.

The present invention will be described in more detail with reference to the following examples, which are not intended to limit the present invention.

### Examples


Human patients with a history of allergy were divided into 5 groups and treated in one eye with eye drop (placebo, 0.01% FK506, 0.03% FK506, 0.06% FK506 or 0.1% FK506), and the other eye with placebo. Each eye drop was administered 4 times per day for seven days and 16 hours after the final instillation, patients were administered allergen-containing eye drops at a concentration predetermined to cause a reaction in the patient. The one hundred patients having a baseline itching score of at least 3, on a scale of 0 to 4, with 4 being most severe, were evaluated. Data of decrease from baseline itching score are presented in Figure 1.

As seen in Figure 1, there was a pronounced dose response, especially at 3 minutes post challenge, when all concentrations were statistically significant versus placebo.

This application is based on application No. 60/402,051 filed in United States of America, the content of which is incorporated herein by reference.

### Claims

1. A method of treating a human patient suffering from ocular allergy, comprising administering to said patient an ophthalmic composition containing from about 0.01% to about 0.1% of macrolide compound.
2. A method according to claim 1 wherein said ocular allergy is allergic conjunctivitis.
3. A method according to claim 1 or 2 wherein said composition contains from about 0.03% to about 0.06% of said macrolide compound.
4. A method according to claim 3 wherein said macrolide compound composition contains about 0.03% of said macrolide compound.
5. A method according to claim 1 wherein said macrolide compound is FK506.
6. A method according to claim 1 wherein said ophthalmic composition is eye drop.
7. A method according to claim 6, wherein said eye drop further contains polyvinyl alcohol.
8. A method according to claim 7, wherein said eye drop contains about 0.03% of said macrolide compound.
9. A method according to claim 8, wherein said eye drop is administered from about one to about 4 times per day.
10. A method according to any of claims 1 to 9, wherein said macrolide compound is a compound having the following formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein adjacent pairs of R<sup>1</sup> and R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup>, and R<sup>5</sup> and R<sup>6</sup> each independently

a) consist of two adjacent hydrogen atoms, wherein R<sup>2</sup> is optionally alkyl, or

b) form another bond optionally between carbon atoms binding with the members of said pairs;

R<sup>7</sup> is hydrogen atom, hydroxy, alkyloxy or protected hydroxy, or may form oxo with R<sup>1</sup>;

R<sup>8</sup> and R<sup>9</sup> each independently show hydrogen atom or hydroxy;

R<sup>10</sup> is hydrogen atom, alkyl, alkyl substituted by one or more hydroxy, alkenyl, alkenyl substituted by one or more hydroxy or alkyl substituted by oxo;

X is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula -CH<sub>2</sub>O-;

Y is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula N-NR<sup>11</sup>R<sup>12</sup> or N-OR<sup>13</sup>;


R<sup>11</sup> and R<sup>12</sup> each independently show hydrogen atom, alkyl, aryl or tosyl;

R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup>, R<sup>22</sup> and R<sup>23</sup> each independently show hydrogen atom or alkyl;

R<sup>24</sup> is an optionally substituted ring that may contain one or more hetero atom(s); and

n is 1 or 2.


11. A method according to claim 10, wherein said macrolide compound has the following structure:



12. An ophthalmic composition for treatment of ocular allergy containing from about 0.01% to about 0.1% of macrolide compound.
13. An ophthalmic composition according to claim 12 wherein said ocular allergy is allergic conjunctivitis.
14. An ophthalmic composition according to claim 12 or 13 which contains from about 0.03% to about 0.06% of said macrolide compound.
15. An ophthalmic composition according to claim 14 which contains about 0.03% of said macrolide compound.
16. An ophthalmic composition according to claim 12 wherein said macrolide compound is FK506.
17. An ophthalmic composition according to claim 12 which is an eye drop.
18. An ophthalmic composition according to claim 17, wherein said eye drop further contains polyvinyl alcohol.
19. An ophthalmic composition according to claim 18, wherein said eye drop contains about 0.03% of said macrolide compound.
20. An ophthalmic composition according to claim 19, wherein said eye drop is administered from about one to about 4 times per day.



21. An ophthalmic composition according to any of claims 12 to 20, wherein said macrolide compound is a compound having the following formula (I) or a pharmaceutically acceptable salt thereof:



wherein adjacent pairs of  $R^1$  and  $R^2$ ,  $R^3$  and  $R^4$ , and  $R^5$  and  $R^6$  each independently

a) consist of two adjacent hydrogen atoms, wherein  $R^2$  is optionally alkyl, or

b) form another bond optionally between carbon atoms binding with the members of said pairs;

$R^7$  is hydrogen atom, hydroxy, alkyloxy or protected hydroxy, or may form oxo with  $R^1$ ;

$R^8$  and  $R^9$  each independently show hydrogen atom or hydroxy;

$R^{10}$  is hydrogen atom, alkyl, alkyl substituted by one or more hydroxy, alkenyl, alkenyl substituted by one or more hydroxy or alkyl substituted by oxo;

X is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $-\text{CH}_2\text{O}-$ ;

Y is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $\text{N}-\text{NR}^{11}\text{R}^{12}$  or  $\text{N}-\text{OR}^{13}$ ;


$R^{11}$  and  $R^{12}$  each independently show hydrogen atom, alkyl, aryl or tosyl;

$R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$ ,  $R^{19}$ ,  $R^{22}$  and  $R^{23}$  each independently show hydrogen atom or alkyl;

$R^{24}$  is an optionally substituted ring that may contain one or more hetero atom(s); and


$n$  is 1 or 2.

22. An ophthalmic composition according to claim 21, wherein said macrolide compound has the following structure:



23. A use of macrolide compound for manufacturing an ophthalmic composition for treatment of ocular allergy, wherein said composition contains from about 0.01% to about 0.1% of said macrolide compound.
24. A use according to claim 23 wherein said ocular allergy is allergic conjunctivitis.
25. A use according to claim 23 or 24 wherein said composition contains from about 0.03% to about 0.06% of said macrolide compound.
26. A use according to claim 25 wherein said composition contains about 0.03% of said macrolide compound.
27. A use according to claim 23 wherein said macrolide compound is FK506.
28. A use according to claim 23 wherein said ophthalmic composition is an eye drop.
29. A use according to claim 28, wherein said eye drop further comprises polyvinyl alcohol.
30. A use according to claim 29, wherein said eye drop contains about 0.03% of said macrolide compound.
31. A use according to claim 30, wherein said eye drop is administered from about one to about 4 times per day.

32. A use according to any of claims 23 to 31, wherein said macrolide compound is a compound having the following formula (I) or a pharmaceutically acceptable salt thereof:



wherein adjacent pairs of  $R^1$  and  $R^2$ ,  $R^3$  and  $R^4$ , and  $R^5$  and  $R^6$  each independently

- a) consist of two adjacent hydrogen atoms, wherein  $R^2$  is optionally alkyl, or
- b) form another bond optionally between carbon atoms binding with the members of said pairs;

$R^7$  is hydrogen atom, hydroxy, alkyloxy or protected hydroxy, or may form oxo with  $R^1$ ;

$R^8$  and  $R^9$  each independently show hydrogen atom or hydroxy;

$R^{10}$  is hydrogen atom, alkyl, alkyl substituted by one or more hydroxy, alkenyl, alkenyl substituted by one or more hydroxy or alkyl substituted by oxo;

X is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $-\text{CH}_2\text{O}-$ ;

Y is oxo, (hydrogen atom, hydroxy), (hydrogen atom, hydrogen atom), or a group of the formula  $\text{N}-\text{NR}^{11}\text{R}^{12}$  or  $\text{N}-\text{OR}^{13}$ ;


$R^{11}$  and  $R^{12}$  each independently show hydrogen atom, alkyl, aryl or tosyl;

$R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$ ,  $R^{19}$ ,  $R^{22}$  and  $R^{23}$  each independently show hydrogen atom or alkyl;


R<sup>24</sup> is an optionally substituted ring that may contain one or more hetero atom(s); and

$n$  is 1 or 2.

33. A use according to claim 32, wherein said macrolide compound has the following structure:



34. A commercial package comprising the ophthalmic composition of any of claims 12 to 22 and a written matter associated therewith, the written matter stating that the composition can or should be used for allergic conjunctivitis.

**Figure 1**

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 03/10105

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K31/436 A61P27/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 03 043650 A (SUCAMPO AG ;UENO RYUJI (US)) 30 May 2003 (2003-05-30) claims 1,4,6,12,15,17,18,31,33,34; table 1	1-34
X	WO 92 19278 A (KURUME UNIVERSITY) 12 November 1992 (1992-11-12) page 1, line 9-11 claims 1,2,5,8 page 2, line 1-4 page 8, line 1-15 page 11, line 1,2 examples 1,2	1-34

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

16 October 2003

Date of mailing of the international search report

24/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Veronese, A

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 03/10105

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 085359 A (SUCAMPO AG ;UENO RYUJI (US)) 31 October 2002 (2002-10-31)  page 15, line 10-13 page 16, line 16-19 page 18, line 14-21 claims 1,2,4,14; examples; tables ---	12,13, 16,17, 20-23
X	EP 0 484 936 A (FUJISAWA PHARMACEUTICAL CO) 13 May 1992 (1992-05-13)  page 6, line 28,29 Prescription 4-6 claims 1,4 ---	12,13, 16,17, 20-22
X	EP 0 406 791 A (FUJISAWA PHARMACEUTICAL CO) 9 January 1991 (1991-01-09)  page 6, line 51 claims 1-3 ---	12,13, 16,17, 20-22
X	WO 00 66122 A (UENO SEIYAKU OYO KENKYUJO KK ;UENO RYUJI (US)) 9 November 2000 (2000-11-09) page 1, line 15,16 example 1 ---	1-34
A	"FUJISAWA Protopic (tacrolimus) FOR DERMATOLOGIC USE ONLY NOT FOR OPHTHALMIC use" INTERNET, December 2000 (2000-12), XP002230797 Retrieved from the Internet: <URL:http://www.protopic.com/img/protopic_pi.pdf> 'retrieved on 2003-02-05! the whole document -----	1-34

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/JP 03/10105

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 1-11 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 03/10105

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03043650	A	30-05-2003	WO 03043650 A1	30-05-2003
WO 9219278	A	12-11-1992	AT 198708 T	15-02-2001
			CA 2102241 A1	27-10-1992
			DE 69231644 D1	22-02-2001
			DE 69231644 T2	23-05-2001
			DK 581959 T3	29-01-2001
			EP 0581959 A1	09-02-1994
			ES 2154262 T3	01-04-2001
			GR 3035269 T3	30-04-2001
			WO 9219278 A1	12-11-1992
			JP 3158437 B2	23-04-2001
			JP 7500570 T	19-01-1995
			KR 237715 B1	01-02-2000
			US 5514686 A	07-05-1996
WO 02085359	A	31-10-2002	WO 02085359 A1	31-10-2002
			US 2002187998 A1	12-12-2002
EP 0484936	A	13-05-1992	AT 112486 T	15-10-1994
			AU 653556 B2	06-10-1994
			AU 8709991 A	14-05-1992
			CA 2054983 A1	09-05-1992
			CN 1061907 A ,B	17-06-1992
			DE 69104460 D1	10-11-1994
			DE 69104460 T2	09-02-1995
			DK 484936 T3	27-03-1995
			EP 0484936 A1	13-05-1992
			ES 2061149 T3	01-12-1994
			HU 60925 A2	30-11-1992
			IE 913861 A1	20-05-1992
			IL 100011 A	08-12-1995
			JP 2581359 B2	12-02-1997
			JP 5155770 A	22-06-1993
			KR 211085 B1	15-07-1999
			PT 99461 A ,B	30-10-1992
			US 5368865 A	29-11-1994
			US 5496564 A	05-03-1996
			RU 2079304 C1	20-05-1997
			ZA 9108846 A	26-08-1992
EP 0406791	A	09-01-1991	AT 117897 T	15-02-1995
			AU 635286 B2	18-03-1993
			AU 5864290 A	24-01-1991
			CA 2020431 A1	06-01-1991
			CN 1048496 A ,B	16-01-1991
			DE 69016515 D1	16-03-1995
			DE 69016515 T2	08-06-1995
			DK 406791 T3	27-03-1995
			EP 0406791 A2	09-01-1991
			ES 2066915 T3	16-03-1995
			GR 3014946 T3	31-05-1995
			HK 27097 A	06-03-1997
			IE 902413 A1	19-06-1991
			IL 94971 A	08-12-1995
			JP 2536248 B2	18-09-1996
			JP 3128320 A	31-05-1991
			KR 163035 B1	01-12-1998

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 03/10105

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0406791	A	US 5770607 A	23-06-1998
WO 0066122	A	09-11-2000	
		AU 4142900 A	17-11-2000
		BR 0011225 A	19-03-2002
		CA 2372448 A1	09-11-2000
		CN 1353608 T	12-06-2002
		CZ 20013769 A3	13-03-2002
		EP 1173177 A1	23-01-2002
		HU 0200864 A2	29-07-2002
		WO 0066122 A1	09-11-2000
		JP 2002543132 T	17-12-2002
		NO 20015288 A	29-10-2001
		TR 200103119 T2	22-04-2002
		US 2003130301 A1	10-07-2003
		ZA 200108905 A	29-10-2002